

Table 3.1A Perceived Risk and Availability of Drugs, by Age Group: Numbers in Thousands, 2002

| Risk/Availability ¹ | Total | AGE GROUP (Years) | | |
|---|---------|-------------------|--------|-------------|
| | | 12-17 | 18-25 | 26 or Older |
| PERCEPTIONS OF GREAT RISK | | | | |
| Cigarettes | | | | |
| Smoke one or more packs per day | 166,124 | 15,493 | 20,152 | 130,478 |
| Marijuana | | | | |
| Smoke once a month | 88,596 | 7,918 | 7,252 | 73,427 |
| Smoke once or twice a week | 118,697 | 12,614 | 10,947 | 95,135 |
| Cocaine | | | | |
| Use once a month | 165,536 | 12,296 | 19,683 | 133,558 |
| Use once or twice a week | 207,016 | 19,463 | 26,794 | 160,759 |
| Heroin | | | | |
| Try once or twice | 190,861 | 14,211 | 23,917 | 152,733 |
| Use once or twice a week | 217,455 | 20,073 | 28,723 | 168,658 |
| LSD | | | | |
| Try once or twice | 169,629 | 12,499 | 19,046 | 138,085 |
| Use once or twice a week | 204,413 | 18,156 | 25,890 | 160,367 |
| Alcohol | | | | |
| Four or five drinks nearly every day | 161,928 | 15,288 | 19,201 | 127,439 |
| Five or more drinks once or twice a week | 98,785 | 9,379 | 10,248 | 79,158 |
| PERCEIVED AVAILABILITY | | | | |
| Fairly or very easy to obtain | | | | |
| Marijuana | 130,549 | 13,299 | 23,718 | 93,532 |
| Cocaine | 70,675 | 5,981 | 12,176 | 52,518 |
| Crack | 65,373 | 6,324 | 9,805 | 49,244 |
| Heroin | 40,380 | 3,765 | 6,062 | 30,554 |
| LSD | 44,659 | 4,562 | 8,634 | 31,463 |
| Approached by someone selling drugs in the past month | 18,152 | 4,121 | 5,894 | 8,137 |

*Low precision; no estimate reported.

¹ Respondents with missing data were excluded.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002.

Table 3.1B Perceived Risk and Availability of Drugs, by Age Group: Percentages, 2002

| Risk/Availability ¹ | Total | AGE GROUP (Years) | | |
|---|-------|-------------------|-------|-------------|
| | | 12-17 | 18-25 | 26 or Older |
| PERCEPTIONS OF GREAT RISK | | | | |
| Cigarettes | | | | |
| Smoke one or more packs per day | 71.1 | 63.1 | 65.2 | 73.3 |
| Marijuana | | | | |
| Smoke once a month | 38.3 | 32.4 | 23.5 | 41.7 |
| Smoke once or twice a week | 51.3 | 51.5 | 35.5 | 54.1 |
| Cocaine | | | | |
| Use once a month | 71.5 | 50.5 | 64.1 | 75.7 |
| Use once or twice a week | 89.4 | 79.8 | 87.2 | 91.1 |
| Heroin | | | | |
| Try once or twice | 82.4 | 58.5 | 78.0 | 86.5 |
| Use once or twice a week | 93.9 | 82.5 | 93.6 | 95.5 |
| LSD | | | | |
| Try once or twice | 73.7 | 52.6 | 62.4 | 78.5 |
| Use once or twice a week | 88.8 | 76.2 | 84.8 | 91.2 |
| Alcohol | | | | |
| Four or five drinks nearly every day | 69.4 | 62.2 | 62.1 | 71.7 |
| Five or more drinks once or twice a week | 42.3 | 38.2 | 33.2 | 44.5 |
| PERCEIVED AVAILABILITY | | | | |
| Fairly or very easy to obtain | | | | |
| Marijuana | 58.0 | 55.0 | 77.4 | 54.9 |
| Cocaine | 31.8 | 25.0 | 40.5 | 31.3 |
| Crack | 29.5 | 26.5 | 32.7 | 29.4 |
| Heroin | 18.3 | 15.8 | 20.3 | 18.3 |
| LSD | 20.3 | 19.4 | 28.9 | 18.9 |
| Approached by someone selling drugs in the past month | 7.7 | 16.7 | 19.0 | 4.5 |

*Low precision; no estimate reported.

¹ Respondents with missing data were excluded.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002.

Table 3.2A Perceived Great Risk of Smoking One or More Packs of Cigarettes Per Day, by Demographic Characteristics (I), and Selected Drug Use, by Risk Perceptions of Smoking One or More Packs of Cigarettes Per Day (II) among Persons Aged 12 to 17: Numbers in Thousands, 2002

| (I) Demographic Characteristic | Smoke One or More Packs of Cigarettes Per Day ¹ | (II) Drug Use | Smoke One or More Packs of Cigarettes Per Day ¹ | |
|---|--|--|--|-------------------------|
| | Great Risk | | Great Risk | Moderate/Slight/No Risk |
| TOTAL | 15,493 | PAST MONTH | | |
| AGE | | Any Illicit Drug ² | 1,560 | 1,312 |
| 12-13 | 5,042 | Marijuana | 1,086 | 937 |
| 14-15 | 5,266 | Any Illicit Drug Other Than Marijuana ² | 749 | 666 |
| 16-17 | 5,185 | Cigarettes | 1,591 | 1,609 |
| GENDER | | Alcohol | 2,456 | 1,892 |
| Male | 7,433 | Binge Alcohol ³ | 1,424 | 1,211 |
| Female | 8,060 | PAST YEAR | | |
| HISPANIC ORIGIN AND RACE | | Any Illicit Drug ² | 3,051 | 2,430 |
| Not Hispanic or Latino | 13,074 | Marijuana | 2,164 | 1,740 |
| White | 9,741 | Any Illicit Drug Other Than Marijuana ² | 1,793 | 1,531 |
| Black or African American | 2,275 | Cigarettes | 2,673 | 2,346 |
| American Indian or Alaska Native | 96 | Alcohol | 5,051 | 3,487 |
| Native Hawaiian or Other Pacific Islander | * | LIFETIME | | |
| Asian | 648 | Any Illicit Drug ² | 4,339 | 3,274 |
| Two or More Races | 266 | Marijuana | 2,919 | 2,178 |
| Hispanic or Latino | 2,419 | Any Illicit Drug Other Than Marijuana ² | 2,911 | 2,349 |
| COUNTY TYPE | | Cigarettes | 4,685 | 3,529 |
| Large Metro | 7,946 | Alcohol | 6,464 | 4,249 |
| Small Metro | 4,554 | | | |
| Nonmetro | 2,993 | | | |

*Low precision; no estimate reported.

¹ Respondents with missing data were excluded.

² Any Illicit Drug includes marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically. Any Illicit Drug Other Than Marijuana includes cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002.

Table 3.2B Perceived Great Risk of Smoking One or More Packs of Cigarettes Per Day, by Demographic Characteristics (I), and Selected Drug Use, by Risk Perceptions of Smoking One or More Packs of Cigarettes Per Day (II) among Persons Aged 12 to 17: Percentages, 2002

| (I) Demographic Characteristic | Smoke One or More Packs of Cigarettes Per Day ¹ | (II) Drug Use | Smoke One or More Packs of Cigarettes Per Day ¹ | |
|---|--|--|--|-------------------------|
| | Great Risk | | Great Risk | Moderate/Slight/No Risk |
| TOTAL | 63.1 | PAST MONTH | | |
| AGE | | Any Illicit Drug ² | 10.1 | 14.5 |
| 12-13 | 61.8 | Marijuana | 7.0 | 10.4 |
| 14-15 | 62.6 | Any Illicit Drug Other Than Marijuana ² | 4.8 | 7.4 |
| 16-17 | 65.1 | Cigarettes | 10.3 | 17.8 |
| GENDER | | Alcohol | 15.9 | 20.9 |
| Male | 59.2 | Binge Alcohol ³ | 9.2 | 13.4 |
| Female | 67.3 | PAST YEAR | | |
| HISPANIC ORIGIN AND RACE | | Any Illicit Drug ² | 19.7 | 26.9 |
| Not Hispanic or Latino | 63.3 | Marijuana | 14.0 | 19.2 |
| White | 63.2 | Any Illicit Drug Other Than Marijuana ² | 11.6 | 16.9 |
| Black or African American | 63.5 | Cigarettes | 17.3 | 25.9 |
| American Indian or Alaska Native | 51.3 | Alcohol | 32.6 | 38.6 |
| Native Hawaiian or Other Pacific Islander | * | LIFETIME | | |
| Asian | 65.2 | Any Illicit Drug ² | 28.0 | 36.2 |
| Two or More Races | 68.0 | Marijuana | 18.8 | 24.1 |
| Hispanic or Latino | 62.4 | Any Illicit Drug Other Than Marijuana ² | 18.8 | 26.0 |
| COUNTY TYPE | | Cigarettes | 30.2 | 39.0 |
| Large Metro | 64.6 | Alcohol | 41.7 | 47.0 |
| Small Metro | 62.3 | | | |
| Nonmetro | 60.8 | | | |

*Low precision; no estimate reported.

¹ Respondents with missing data were excluded.

² Any Illicit Drug includes marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically. Any Illicit Drug Other Than Marijuana includes cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002.

Table 3.3A Perceived Great Risk of Smoking Marijuana Once a Month, by Demographic Characteristics (I), and Selected Drug Use, by Risk Perceptions of Smoking Marijuana Once a Month (II) among Persons Aged 12 to 17: Numbers in Thousands, 2002

| (I) Demographic Characteristic | Smoke Marijuana Once a Month ¹ | (II) Drug Use | Smoke Marijuana Once a Month ¹ | |
|---|---|--|---|-------------------------|
| | Great Risk | | Great Risk | Moderate/Slight/No Risk |
| TOTAL | 7,918 | PAST MONTH | | |
| AGE | | Any Illicit Drug ² | 383 | 2,485 |
| 12-13 | 3,411 | Marijuana | 150 | 1,870 |
| 14-15 | 2,586 | Any Illicit Drug Other Than Marijuana ² | 262 | 1,152 |
| 16-17 | 1,920 | Cigarettes | 475 | 2,719 |
| GENDER | | Alcohol | 747 | 3,593 |
| Male | 3,836 | Binge Alcohol ³ | 390 | 2,238 |
| Female | 4,081 | PAST YEAR | | |
| HISPANIC ORIGIN AND RACE | | Any Illicit Drug ² | 834 | 4,637 |
| Not Hispanic or Latino | 6,572 | Marijuana | 375 | 3,523 |
| White | 4,829 | Any Illicit Drug Other Than Marijuana ² | 592 | 2,728 |
| Black or African American | 1,229 | Cigarettes | 873 | 4,142 |
| American Indian or Alaska Native | 54 | Alcohol | 1,718 | 6,789 |
| Native Hawaiian or Other Pacific Islander | * | LIFETIME | | |
| Asian | 311 | Any Illicit Drug ² | 1,399 | 6,195 |
| Two or More Races | 126 | Marijuana | 620 | 4,471 |
| Hispanic or Latino | 1,346 | Any Illicit Drug Other Than Marijuana ² | 1,067 | 4,180 |
| COUNTY TYPE | | Cigarettes | 1,710 | 6,488 |
| Large Metro | 3,844 | Alcohol | 2,379 | 8,305 |
| Small Metro | 2,259 | | | |
| Nonmetro | 1,815 | | | |

*Low precision; no estimate reported.

¹ Respondents with missing data were excluded.

² Any Illicit Drug includes marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically. Any Illicit Drug Other Than Marijuana includes cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002.

Table 3.3B Perceived Great Risk of Smoking Marijuana Once a Month, by Demographic Characteristics (I), and Selected Drug Use, by Risk Perceptions of Smoking Marijuana Once a Month (II) among Persons Aged 12 to 17: Percentages, 2002

| (I) Demographic Characteristic | Smoke Marijuana Once a Month ¹ | (II) Drug Use | Smoke Marijuana Once a Month ¹ | |
|---|---|--|---|-------------------------|
| | Great Risk | | Great Risk | Moderate/Slight/No Risk |
| TOTAL | 32.4 | PAST MONTH | | |
| AGE | | Any Illicit Drug ² | 4.8 | 15.0 |
| 12-13 | 42.0 | Marijuana | 1.9 | 11.3 |
| 14-15 | 30.9 | Any Illicit Drug Other Than Marijuana ² | 3.3 | 7.0 |
| 16-17 | 24.1 | Cigarettes | 6.0 | 16.5 |
| GENDER | | Alcohol | 9.4 | 21.7 |
| Male | 30.7 | Binge Alcohol ³ | 4.9 | 13.5 |
| Female | 34.1 | PAST YEAR | | |
| HISPANIC ORIGIN AND RACE | | Any Illicit Drug ² | 10.5 | 28.1 |
| Not Hispanic or Latino | 32.0 | Marijuana | 4.7 | 21.3 |
| White | 31.4 | Any Illicit Drug Other Than Marijuana ² | 7.5 | 16.5 |
| Black or African American | 34.4 | Cigarettes | 11.0 | 25.1 |
| American Indian or Alaska Native | 28.9 | Alcohol | 21.7 | 41.1 |
| Native Hawaiian or Other Pacific Islander | * | LIFETIME | | |
| Asian | 31.8 | Any Illicit Drug ² | 17.7 | 37.5 |
| Two or More Races | 33.0 | Marijuana | 7.8 | 27.1 |
| Hispanic or Latino | 34.8 | Any Illicit Drug Other Than Marijuana ² | 13.5 | 25.3 |
| COUNTY TYPE | | Cigarettes | 21.6 | 39.3 |
| Large Metro | 31.3 | Alcohol | 30.1 | 50.3 |
| Small Metro | 31.0 | | | |
| Nonmetro | 37.1 | | | |

*Low precision; no estimate reported.

¹ Respondents with missing data were excluded.

² Any Illicit Drug includes marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically. Any Illicit Drug Other Than Marijuana includes cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002.

Table 3.4A Perceived Great Risk of Smoking Marijuana Once or Twice a Week, by Demographic Characteristics (I), and Selected Drug Use, by Risk Perceptions of Smoking Marijuana Once or Twice a Week (II) among Persons Aged 12 to 17: Numbers in Thousands, 2002

| (I) Demographic Characteristic | Smoke Marijuana Once or Twice a Week ¹ | (II) Drug Use | Smoke Marijuana Once or Twice a Week ¹ | |
|---|---|--|---|-------------------------|
| | Great Risk | | Great Risk | Moderate/Slight/No Risk |
| TOTAL | 12,614 | PAST MONTH | | |
| AGE | | Any Illicit Drug ² | 534 | 2,333 |
| 12-13 | 5,197 | Marijuana | 208 | 1,812 |
| 14-15 | 4,177 | Any Illicit Drug Other Than Marijuana ² | 370 | 1,043 |
| 16-17 | 3,240 | Cigarettes | 734 | 2,454 |
| GENDER | | Alcohol | 1,150 | 3,188 |
| Male | 5,954 | Binge Alcohol ³ | 556 | 2,072 |
| Female | 6,660 | PAST YEAR | | |
| HISPANIC ORIGIN AND RACE | | Any Illicit Drug ² | 1,318 | 4,149 |
| Not Hispanic or Latino | 10,660 | Marijuana | 616 | 3,279 |
| White | 8,184 | Any Illicit Drug Other Than Marijuana ² | 882 | 2,437 |
| Black or African American | 1,629 | Cigarettes | 1,352 | 3,659 |
| American Indian or Alaska Native | 83 | Alcohol | 2,828 | 5,680 |
| Native Hawaiian or Other Pacific Islander | * | LIFETIME | | |
| Asian | 545 | Any Illicit Drug ² | 2,222 | 5,361 |
| Two or More Races | 191 | Marijuana | 1,008 | 4,079 |
| Hispanic or Latino | 1,955 | Any Illicit Drug Other Than Marijuana ² | 1,621 | 3,617 |
| COUNTY TYPE | | Cigarettes | 2,645 | 5,551 |
| Large Metro | 6,157 | Alcohol | 3,857 | 6,828 |
| Small Metro | 3,740 | | | |
| Nonmetro | 2,718 | | | |

*Low precision; no estimate reported.

¹ Respondents with missing data were excluded.

² Any Illicit Drug includes marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically. Any Illicit Drug Other Than Marijuana includes cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002.

Table 3.4B Perceived Great Risk of Smoking Marijuana Once or Twice a Week, by Demographic Characteristics (I), and Selected Drug Use, by Risk Perceptions of Smoking Marijuana Once or Twice a Week (II) among Persons Aged 12 to 17: Percentages, 2002

| (I) Demographic Characteristic | Smoke Marijuana Once or Twice a Week ¹ | (II) Drug Use | Smoke Marijuana Once or Twice a Week ¹ | |
|---|---|--|---|-------------------------|
| | Great Risk | | Great Risk | Moderate/Slight/No Risk |
| TOTAL | 51.5 | PAST MONTH | | |
| AGE | | Any Illicit Drug ² | 4.2 | 19.7 |
| 12-13 | 63.8 | Marijuana | 1.7 | 15.3 |
| 14-15 | 49.8 | Any Illicit Drug Other Than Marijuana ² | 2.9 | 8.8 |
| 16-17 | 40.7 | Cigarettes | 5.8 | 20.7 |
| GENDER | | Alcohol | 9.1 | 26.9 |
| Male | 47.6 | Binge Alcohol ³ | 4.4 | 17.5 |
| Female | 55.7 | PAST YEAR | | |
| HISPANIC ORIGIN AND RACE | | Any Illicit Drug ² | 10.5 | 35.0 |
| Not Hispanic or Latino | 51.8 | Marijuana | 4.9 | 27.6 |
| White | 53.1 | Any Illicit Drug Other Than Marijuana ² | 7.0 | 20.5 |
| Black or African American | 45.6 | Cigarettes | 10.7 | 30.8 |
| American Indian or Alaska Native | 44.3 | Alcohol | 22.4 | 47.9 |
| Native Hawaiian or Other Pacific Islander | * | LIFETIME | | |
| Asian | 55.7 | Any Illicit Drug ² | 17.6 | 45.2 |
| Two or More Races | 49.9 | Marijuana | 8.0 | 34.4 |
| Hispanic or Latino | 50.3 | Any Illicit Drug Other Than Marijuana ² | 12.8 | 30.5 |
| COUNTY TYPE | | Cigarettes | 21.0 | 46.8 |
| Large Metro | 50.2 | Alcohol | 30.6 | 57.5 |
| Small Metro | 51.2 | | | |
| Nonmetro | 55.5 | | | |

*Low precision; no estimate reported.

¹ Respondents with missing data were excluded.

² Any Illicit Drug includes marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically. Any Illicit Drug Other Than Marijuana includes cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002.

Table 3.5A Perceived Great Risk of Using Cocaine Once a Month, by Demographic Characteristics (I), and Selected Drug Use, by Risk Perceptions of Using Cocaine Once a Month (II) among Persons Aged 12 to 17: Numbers in Thousands, 2002

| (I) Demographic Characteristic | Use Cocaine Once a Month ¹ | (II) Drug Use | Use Cocaine Once a Month ¹ | |
|---|---------------------------------------|--|---------------------------------------|-------------------------|
| | Great Risk | | Great Risk | Moderate/Slight/No Risk |
| TOTAL | 12,296 | PAST MONTH | | |
| AGE | | Any Illicit Drug ² | 1,376 | 1,476 |
| 12-13 | 3,461 | Marijuana | 990 | 1,020 |
| 14-15 | 4,135 | Any Illicit Drug Other Than Marijuana ² | 613 | 794 |
| 16-17 | 4,700 | Cigarettes | 1,731 | 1,440 |
| GENDER | | Alcohol | 2,316 | 2,009 |
| Male | 6,316 | Binge Alcohol ³ | 1,365 | 1,254 |
| Female | 5,980 | PAST YEAR | | |
| HISPANIC ORIGIN AND RACE | | Any Illicit Drug ² | 2,764 | 2,669 |
| Not Hispanic or Latino | 10,507 | Marijuana | 2,073 | 1,789 |
| White | 7,805 | Any Illicit Drug Other Than Marijuana ² | 1,452 | 1,860 |
| Black or African American | 1,977 | Cigarettes | 2,684 | 2,285 |
| American Indian or Alaska Native | 84 | Alcohol | 4,582 | 3,888 |
| Native Hawaiian or Other Pacific Islander | * | LIFETIME | | |
| Asian | 432 | Any Illicit Drug ² | 3,905 | 3,626 |
| Two or More Races | 184 | Marijuana | 2,836 | 2,208 |
| Hispanic or Latino | 1,789 | Any Illicit Drug Other Than Marijuana ² | 2,412 | 2,806 |
| COUNTY TYPE | | Cigarettes | 4,470 | 3,668 |
| Large Metro | 5,994 | Alcohol | 5,787 | 4,840 |
| Small Metro | 3,739 | | | |
| Nonmetro | 2,564 | | | |

*Low precision; no estimate reported.

¹ Respondents with missing data were excluded.

² Any Illicit Drug includes marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically. Any Illicit Drug Other Than Marijuana includes cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002.

Table 3.5B Perceived Great Risk of Using Cocaine Once a Month, by Demographic Characteristics (I), and Selected Drug Use, by Risk Perceptions of Using Cocaine Once a Month (II) among Persons Aged 12 to 17: Percentages, 2002

| (I) Demographic Characteristic | Use Cocaine Once a Month ¹ | (II) Drug Use | Use Cocaine Once a Month ¹ | |
|---|---------------------------------------|--|---------------------------------------|-------------------------|
| | Great Risk | | Great Risk | Moderate/Slight/No Risk |
| TOTAL | 50.5 | PAST MONTH | | |
| AGE | | Any Illicit Drug ² | 11.2 | 12.2 |
| 12-13 | 42.8 | Marijuana | 8.1 | 8.5 |
| 14-15 | 49.7 | Any Illicit Drug Other Than Marijuana ² | 5.0 | 6.6 |
| 16-17 | 59.2 | Cigarettes | 14.1 | 11.9 |
| GENDER | | Alcohol | 18.8 | 16.7 |
| Male | 50.7 | Binge Alcohol ³ | 11.1 | 10.4 |
| Female | 50.2 | PAST YEAR | | |
| HISPANIC ORIGIN AND RACE | | Any Illicit Drug ² | 22.5 | 22.1 |
| Not Hispanic or Latino | 51.3 | Marijuana | 16.9 | 14.8 |
| White | 50.9 | Any Illicit Drug Other Than Marijuana ² | 11.8 | 15.4 |
| Black or African American | 55.9 | Cigarettes | 21.8 | 18.9 |
| American Indian or Alaska Native | 45.1 | Alcohol | 37.3 | 32.2 |
| Native Hawaiian or Other Pacific Islander | * | LIFETIME | | |
| Asian | 44.2 | Any Illicit Drug ² | 31.8 | 30.1 |
| Two or More Races | 47.9 | Marijuana | 23.1 | 18.3 |
| Hispanic or Latino | 46.3 | Any Illicit Drug Other Than Marijuana ² | 19.6 | 23.3 |
| COUNTY TYPE | | Cigarettes | 36.4 | 30.4 |
| Large Metro | 49.1 | Alcohol | 47.1 | 40.1 |
| Small Metro | 51.5 | | | |
| Nonmetro | 52.5 | | | |

*Low precision; no estimate reported.

¹ Respondents with missing data were excluded.

² Any Illicit Drug includes marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically. Any Illicit Drug Other Than Marijuana includes cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002.

Table 3.6A Perceived Great Risk of Using Cocaine Once or Twice a Week, by Demographic Characteristics (I), and Selected Drug Use, by Risk Perceptions of Using Cocaine Once or Twice a Week (II) among Persons Aged 12 to 17: Numbers in Thousands, 2002

| (I) Demographic Characteristic | Use Cocaine Once or Twice a Week ¹ | (II) Drug Use | Use Cocaine Once or Twice a Week ¹ | |
|---|---|--|---|-------------------------|
| | Great Risk | | Great Risk | Moderate/Slight/No Risk |
| TOTAL | 19,463 | PAST MONTH | | |
| AGE | | Any Illicit Drug ² | 2,108 | 744 |
| 12-13 | 5,991 | Marijuana | 1,522 | 488 |
| 14-15 | 6,685 | Any Illicit Drug Other Than Marijuana ² | 961 | 447 |
| 16-17 | 6,787 | Cigarettes | 2,508 | 664 |
| GENDER | | Alcohol | 3,480 | 845 |
| Male | 9,967 | Binge Alcohol ³ | 2,042 | 579 |
| Female | 9,496 | PAST YEAR | | |
| HISPANIC ORIGIN AND RACE | | Any Illicit Drug ² | 4,268 | 1,166 |
| Not Hispanic or Latino | 16,611 | Marijuana | 3,099 | 764 |
| White | 12,776 | Any Illicit Drug Other Than Marijuana ² | 2,417 | 896 |
| Black or African American | 2,613 | Cigarettes | 3,988 | 985 |
| American Indian or Alaska Native | 125 | Alcohol | 6,952 | 1,520 |
| Native Hawaiian or Other Pacific Islander | * | LIFETIME | | |
| Asian | 733 | Any Illicit Drug ² | 5,926 | 1,610 |
| Two or More Races | 312 | Marijuana | 4,127 | 917 |
| Hispanic or Latino | 2,852 | Any Illicit Drug Other Than Marijuana ² | 3,886 | 1,337 |
| COUNTY TYPE | | Cigarettes | 6,581 | 1,560 |
| Large Metro | 9,648 | Alcohol | 8,742 | 1,884 |
| Small Metro | 5,845 | | | |
| Nonmetro | 3,969 | | | |

*Low precision; no estimate reported.

¹ Respondents with missing data were excluded.

² Any Illicit Drug includes marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically. Any Illicit Drug Other Than Marijuana includes cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002.

Table 3.6B Perceived Great Risk of Using Cocaine Once or Twice a Week, by Demographic Characteristics (I), and Selected Drug Use, by Risk Perceptions of Using Cocaine Once or Twice a Week (II) among Persons Aged 12 to 17: Percentages, 2002

| (I) Demographic Characteristic | Use Cocaine Once or Twice a Week ¹ | (II) Drug Use | Use Cocaine Once or Twice a Week ¹ | |
|---|---|--|---|-------------------------|
| | Great Risk | | Great Risk | Moderate/Slight/No Risk |
| TOTAL | 79.8 | PAST MONTH | | |
| AGE | | Any Illicit Drug ² | 10.8 | 15.1 |
| 12-13 | 73.8 | Marijuana | 7.8 | 9.9 |
| 14-15 | 80.3 | Any Illicit Drug Other Than Marijuana ² | 4.9 | 9.1 |
| 16-17 | 85.4 | Cigarettes | 12.9 | 13.5 |
| GENDER | | Alcohol | 17.9 | 17.2 |
| Male | 80.0 | Binge Alcohol ³ | 10.5 | 11.8 |
| Female | 79.6 | PAST YEAR | | |
| HISPANIC ORIGIN AND RACE | | Any Illicit Drug ² | 21.9 | 23.7 |
| Not Hispanic or Latino | 81.0 | Marijuana | 15.9 | 15.5 |
| White | 83.2 | Any Illicit Drug Other Than Marijuana ² | 12.4 | 18.2 |
| Black or African American | 74.0 | Cigarettes | 20.5 | 20.0 |
| American Indian or Alaska Native | 67.2 | Alcohol | 35.7 | 30.8 |
| Native Hawaiian or Other Pacific Islander | * | LIFETIME | | |
| Asian | 74.7 | Any Illicit Drug ² | 30.4 | 32.7 |
| Two or More Races | 81.0 | Marijuana | 21.2 | 18.6 |
| Hispanic or Latino | 73.6 | Any Illicit Drug Other Than Marijuana ² | 20.0 | 27.1 |
| COUNTY TYPE | | Cigarettes | 33.8 | 31.7 |
| Large Metro | 78.9 | Alcohol | 44.9 | 38.2 |
| Small Metro | 80.4 | | | |
| Nonmetro | 81.2 | | | |

*Low precision; no estimate reported.

¹ Respondents with missing data were excluded.

² Any Illicit Drug includes marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically. Any Illicit Drug Other Than Marijuana includes cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002.

Table 3.7A Perceived Great Risk of Trying Heroin Once or Twice, by Demographic Characteristics (I), and Selected Drug Use, by Risk Perceptions of Trying Heroin Once or Twice (II) among Persons Aged 12 to 17: Numbers in Thousands, 2002

| (I) Demographic Characteristic | Try Heroin Once or Twice ¹ | (II) Drug Use | Try Heroin Once or Twice ¹ | |
|---|---------------------------------------|--|---------------------------------------|-------------------------|
| | Great Risk | | Great Risk | Moderate/Slight/No Risk |
| TOTAL | 14,211 | PAST MONTH | | |
| AGE | | Any Illicit Drug ² | 1,991 | 847 |
| 12-13 | 3,828 | Marijuana | 1,480 | 516 |
| 14-15 | 4,816 | Any Illicit Drug Other Than Marijuana ² | 924 | 483 |
| 16-17 | 5,567 | Cigarettes | 2,312 | 842 |
| GENDER | | Alcohol | 2,995 | 1,314 |
| Male | 7,339 | Binge Alcohol ³ | 1,903 | 708 |
| Female | 6,873 | PAST YEAR | | |
| HISPANIC ORIGIN AND RACE | | Any Illicit Drug ² | 3,673 | 1,738 |
| Not Hispanic or Latino | 11,930 | Marijuana | 2,824 | 1,022 |
| White | 8,893 | Any Illicit Drug Other Than Marijuana ² | 2,105 | 1,197 |
| Black or African American | 2,161 | Cigarettes | 3,453 | 1,500 |
| American Indian or Alaska Native | 109 | Alcohol | 5,677 | 2,773 |
| Native Hawaiian or Other Pacific Islander | * | LIFETIME | | |
| Asian | 521 | Any Illicit Drug ² | 4,972 | 2,533 |
| Two or More Races | 215 | Marijuana | 3,677 | 1,340 |
| Hispanic or Latino | 2,281 | Any Illicit Drug Other Than Marijuana ² | 3,291 | 1,917 |
| COUNTY TYPE | | Cigarettes | 5,516 | 2,593 |
| Large Metro | 7,039 | Alcohol | 7,062 | 3,534 |
| Small Metro | 4,252 | | | |
| Nonmetro | 2,921 | | | |

*Low precision; no estimate reported.

¹ Respondents with missing data were excluded.

² Any Illicit Drug includes marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically. Any Illicit Drug Other Than Marijuana includes cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002.

Table 3.7B Perceived Great Risk of Trying Heroin Once or Twice, by Demographic Characteristics (I), and Selected Drug Use, by Risk Perceptions of Trying Heroin Once or Twice (II) among Persons Aged 12 to 17: Percentages, 2002

| (I) Demographic Characteristic | Try Heroin Once or Twice ¹ | (II) Drug Use | Try Heroin Once or Twice ¹ | |
|---|---------------------------------------|--|---------------------------------------|-------------------------|
| | Great Risk | | Great Risk | Moderate/Slight/No Risk |
| TOTAL | 58.5 | PAST MONTH | | |
| AGE | | Any Illicit Drug ² | 14.0 | 8.4 |
| 12-13 | 47.3 | Marijuana | 10.4 | 5.1 |
| 14-15 | 58.1 | Any Illicit Drug Other Than Marijuana ² | 6.5 | 4.8 |
| 16-17 | 70.3 | Cigarettes | 16.3 | 8.3 |
| GENDER | | Alcohol | 21.1 | 13.0 |
| Male | 59.2 | Binge Alcohol ³ | 13.4 | 7.0 |
| Female | 57.7 | PAST YEAR | | |
| HISPANIC ORIGIN AND RACE | | Any Illicit Drug ² | 25.8 | 17.2 |
| Not Hispanic or Latino | 58.4 | Marijuana | 19.9 | 10.1 |
| White | 58.1 | Any Illicit Drug Other Than Marijuana ² | 14.8 | 11.9 |
| Black or African American | 61.2 | Cigarettes | 24.3 | 14.9 |
| American Indian or Alaska Native | 59.0 | Alcohol | 39.9 | 27.5 |
| Native Hawaiian or Other Pacific Islander | * | LIFETIME | | |
| Asian | 53.8 | Any Illicit Drug ² | 35.0 | 25.1 |
| Two or More Races | 55.7 | Marijuana | 25.9 | 13.3 |
| Hispanic or Latino | 59.1 | Any Illicit Drug Other Than Marijuana ² | 23.2 | 19.0 |
| COUNTY TYPE | | Cigarettes | 38.8 | 25.7 |
| Large Metro | 57.8 | Alcohol | 49.7 | 35.0 |
| Small Metro | 58.7 | | | |
| Nonmetro | 60.0 | | | |

*Low precision; no estimate reported.

¹ Respondents with missing data were excluded.

² Any Illicit Drug includes marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically. Any Illicit Drug Other Than Marijuana includes cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002.

Table 3.8A Perceived Great Risk of Using Heroin Once or Twice a Week, by Demographic Characteristics (I), and Selected Drug Use, by Risk Perceptions of Using Heroin Once or Twice a Week (II) among Persons Aged 12 to 17: Numbers in Thousands, 2002

| (I) Demographic Characteristic | Use Heroin Once or Twice a Week ¹ | (II) Drug Use | Use Heroin Once or Twice a Week ¹ | |
|---|--|--|--|-------------------------|
| | Great Risk | | Great Risk | Moderate/Slight/No Risk |
| TOTAL | 20,073 | PAST MONTH | | |
| AGE | | Any Illicit Drug ² | 2,472 | 372 |
| 12-13 | 6,000 | Marijuana | 1,829 | 171 |
| 14-15 | 6,915 | Any Illicit Drug Other Than Marijuana ² | 1,160 | 248 |
| 16-17 | 7,158 | Cigarettes | 2,829 | 333 |
| GENDER | | Alcohol | 3,851 | 465 |
| Male | 10,238 | Binge Alcohol ³ | 2,354 | 261 |
| Female | 9,835 | PAST YEAR | | |
| HISPANIC ORIGIN AND RACE | | Any Illicit Drug ² | 4,692 | 726 |
| Not Hispanic or Latino | 17,055 | Marijuana | 3,515 | 336 |
| White | 13,185 | Any Illicit Drug Other Than Marijuana ² | 2,769 | 535 |
| Black or African American | 2,611 | Cigarettes | 4,394 | 565 |
| American Indian or Alaska Native | 147 | Alcohol | 7,475 | 978 |
| Native Hawaiian or Other Pacific Islander | * | LIFETIME | | |
| Asian | 752 | Any Illicit Drug ² | 6,393 | 1,121 |
| Two or More Races | 314 | Marijuana | 4,563 | 464 |
| Hispanic or Latino | 3,018 | Any Illicit Drug Other Than Marijuana ² | 4,305 | 906 |
| COUNTY TYPE | | Cigarettes | 7,081 | 1,032 |
| Large Metro | 10,007 | Alcohol | 9,274 | 1,327 |
| Small Metro | 5,990 | | | |
| Nonmetro | 4,077 | | | |

*Low precision; no estimate reported.

¹ Respondents with missing data were excluded.

² Any Illicit Drug includes marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically. Any Illicit Drug Other Than Marijuana includes cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002.

Table 3.8B Perceived Great Risk of Using Heroin Once or Twice a Week, by Demographic Characteristics (I), and Selected Drug Use, by Risk Perceptions of Using Heroin Once or Twice a Week (II) among Persons Aged 12 to 17: Percentages, 2002

| (I) Demographic Characteristic | Use Heroin Once or Twice a Week ¹ | (II) Drug Use | Use Heroin Once or Twice a Week ¹ | |
|---|--|--|--|-------------------------|
| | Great Risk | | Great Risk | Moderate/Slight/No Risk |
| TOTAL | 82.5 | PAST MONTH | | |
| AGE | | Any Illicit Drug ² | 12.3 | 8.8 |
| 12-13 | 74.1 | Marijuana | 9.1 | 4.0 |
| 14-15 | 83.4 | Any Illicit Drug Other Than Marijuana ² | 5.8 | 5.8 |
| 16-17 | 90.3 | Cigarettes | 14.1 | 7.8 |
| GENDER | | Alcohol | 19.2 | 10.9 |
| Male | 82.6 | Binge Alcohol ³ | 11.7 | 6.1 |
| Female | 82.5 | PAST YEAR | | |
| HISPANIC ORIGIN AND RACE | | Any Illicit Drug ² | 23.4 | 17.1 |
| Not Hispanic or Latino | 83.3 | Marijuana | 17.5 | 7.9 |
| White | 86.1 | Any Illicit Drug Other Than Marijuana ² | 13.8 | 12.6 |
| Black or African American | 73.9 | Cigarettes | 21.9 | 13.3 |
| American Indian or Alaska Native | 79.4 | Alcohol | 37.2 | 23.0 |
| Native Hawaiian or Other Pacific Islander | * | LIFETIME | | |
| Asian | 77.1 | Any Illicit Drug ² | 31.9 | 26.4 |
| Two or More Races | 81.4 | Marijuana | 22.7 | 10.9 |
| Hispanic or Latino | 78.2 | Any Illicit Drug Other Than Marijuana ² | 21.4 | 21.3 |
| COUNTY TYPE | | Cigarettes | 35.3 | 24.3 |
| Large Metro | 82.0 | Alcohol | 46.2 | 31.2 |
| Small Metro | 82.6 | | | |
| Nonmetro | 83.7 | | | |

*Low precision; no estimate reported.

¹ Respondents with missing data were excluded.

² Any Illicit Drug includes marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically. Any Illicit Drug Other Than Marijuana includes cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002.

Table 3.9A Perceived Great Risk of Trying LSD Once or Twice, by Demographic Characteristics (I), and Selected Drug Use, by Risk Perceptions of Trying LSD Once or Twice (II) among Persons Aged 12 to 17: Numbers in Thousands, 2002

| (I) Demographic Characteristic | Try LSD Once or Twice ¹ | | (II) Drug Use | Try LSD Once or Twice ¹ | |
|---|------------------------------------|--|--|------------------------------------|-------------------------|
| | Great Risk | | | Great Risk | Moderate/Slight/No Risk |
| TOTAL | 12,499 | | PAST MONTH | | |
| AGE | | | Any Illicit Drug ² | 1,309 | 1,480 |
| 12-13 | 3,601 | | Marijuana | 911 | 1,051 |
| 14-15 | 4,252 | | Any Illicit Drug Other Than Marijuana ² | 617 | 764 |
| 16-17 | 4,646 | | Cigarettes | 1,697 | 1,412 |
| GENDER | | | Alcohol | 2,248 | 2,012 |
| Male | 6,323 | | Binge Alcohol ³ | 1,334 | 1,254 |
| Female | 6,175 | | PAST YEAR | | |
| HISPANIC ORIGIN AND RACE | | | Any Illicit Drug ² | 2,603 | 2,721 |
| Not Hispanic or Latino | 10,480 | | Marijuana | 1,905 | 1,890 |
| White | 7,736 | | Any Illicit Drug Other Than Marijuana ² | 1,450 | 1,798 |
| Black or African American | 1,983 | | Cigarettes | 2,633 | 2,256 |
| American Indian or Alaska Native | 92 | | Alcohol | 4,470 | 3,854 |
| Native Hawaiian or Other Pacific Islander | * | | LIFETIME | | |
| Asian | 438 | | Any Illicit Drug ² | 3,760 | 3,623 |
| Two or More Races | 189 | | Marijuana | 2,645 | 2,313 |
| Hispanic or Latino | 2,019 | | Any Illicit Drug Other Than Marijuana ² | 2,420 | 2,695 |
| COUNTY TYPE | | | Cigarettes | 4,371 | 3,617 |
| Large Metro | 6,155 | | Alcohol | 5,687 | 4,749 |
| Small Metro | 3,722 | | | | |
| Nonmetro | 2,622 | | | | |

*Low precision; no estimate reported.

¹ Respondents with missing data were excluded.

² Any Illicit Drug includes marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically. Any Illicit Drug Other Than Marijuana includes cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002.

Table 3.9B Perceived Great Risk of Trying LSD Once or Twice, by Demographic Characteristics (I), and Selected Drug Use, by Risk Perceptions of Trying LSD Once or Twice (II) among Persons Aged 12 to 17: Percentages, 2002

| (I) Demographic Characteristic | Try LSD Once or Twice ¹ | | (II) Drug Use | Try LSD Once or Twice ¹ | |
|---|------------------------------------|--|--|------------------------------------|-------------------------|
| | Great Risk | | | Great Risk | Moderate/Slight/No Risk |
| TOTAL | 52.6 | | PAST MONTH | | |
| AGE | | | Any Illicit Drug ² | 10.5 | 13.1 |
| 12-13 | 46.2 | | Marijuana | 7.3 | 9.3 |
| 14-15 | 52.3 | | Any Illicit Drug Other Than Marijuana ² | 4.9 | 6.8 |
| 16-17 | 59.2 | | Cigarettes | 13.6 | 12.5 |
| GENDER | | | Alcohol | 18.0 | 17.8 |
| Male | 52.1 | | Binge Alcohol ³ | 10.7 | 11.1 |
| Female | 53.0 | | PAST YEAR | | |
| HISPANIC ORIGIN AND RACE | | | Any Illicit Drug ² | 20.8 | 24.1 |
| Not Hispanic or Latino | 52.4 | | Marijuana | 15.2 | 16.8 |
| White | 51.6 | | Any Illicit Drug Other Than Marijuana ² | 11.6 | 15.9 |
| Black or African American | 57.6 | | Cigarettes | 21.1 | 20.0 |
| American Indian or Alaska Native | 50.0 | | Alcohol | 35.8 | 34.2 |
| Native Hawaiian or Other Pacific Islander | * | | LIFETIME | | |
| Asian | 46.8 | | Any Illicit Drug ² | 30.1 | 32.1 |
| Two or More Races | 50.2 | | Marijuana | 21.2 | 20.5 |
| Hispanic or Latino | 53.3 | | Any Illicit Drug Other Than Marijuana ² | 19.4 | 23.9 |
| COUNTY TYPE | | | Cigarettes | 35.0 | 32.1 |
| Large Metro | 51.6 | | Alcohol | 45.5 | 42.1 |
| Small Metro | 52.5 | | | | |
| Nonmetro | 54.9 | | | | |

*Low precision; no estimate reported.

¹ Respondents with missing data were excluded.

² Any Illicit Drug includes marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically. Any Illicit Drug Other Than Marijuana includes cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002.

Table 3.10A Perceived Great Risk of Using LSD Once or Twice a Week, by Demographic Characteristics (I), and Selected Drug Use, by Risk Perceptions of Using LSD Once or Twice a Week (II) among Persons Aged 12 to 17: Numbers in Thousands, 2002

| (I) Demographic Characteristic | Use LSD Once or Twice a Week ¹ | (II) Drug Use | Use LSD Once or Twice a Week ¹ | |
|---|---|--|---|-------------------------|
| | Great Risk | | Great Risk | Moderate/Slight/No Risk |
| TOTAL | 18,156 | PAST MONTH | | |
| AGE | | Any Illicit Drug ² | 2,028 | 769 |
| 12-13 | 5,445 | Marijuana | 1,463 | 509 |
| 14-15 | 6,238 | Any Illicit Drug Other Than Marijuana ² | 941 | 446 |
| 16-17 | 6,473 | Cigarettes | 2,393 | 716 |
| GENDER | | Alcohol | 3,326 | 934 |
| Male | 9,130 | Binge Alcohol ³ | 1,968 | 626 |
| Female | 9,026 | PAST YEAR | | |
| HISPANIC ORIGIN AND RACE | | Any Illicit Drug ² | 3,992 | 1,337 |
| Not Hispanic or Latino | 15,456 | Marijuana | 2,924 | 879 |
| White | 11,930 | Any Illicit Drug Other Than Marijuana ² | 2,303 | 949 |
| Black or African American | 2,409 | Cigarettes | 3,772 | 1,113 |
| American Indian or Alaska Native | 131 | Alcohol | 6,582 | 1,746 |
| Native Hawaiian or Other Pacific Islander | * | LIFETIME | | |
| Asian | 653 | Any Illicit Drug ² | 5,561 | 1,817 |
| Two or More Races | 282 | Marijuana | 3,877 | 1,086 |
| Hispanic or Latino | 2,700 | Any Illicit Drug Other Than Marijuana ² | 3,691 | 1,424 |
| COUNTY TYPE | | Cigarettes | 6,182 | 1,795 |
| Large Metro | 8,999 | Alcohol | 8,222 | 2,217 |
| Small Metro | 5,432 | | | |
| Nonmetro | 3,725 | | | |

*Low precision; no estimate reported.

¹ Respondents with missing data were excluded.

² Any Illicit Drug includes marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically. Any Illicit Drug Other Than Marijuana includes cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002.

Table 3.10B Perceived Great Risk of Using LSD Once or Twice a Week, by Demographic Characteristics (I), and Selected Drug Use, by Risk Perceptions of Using LSD Once or Twice a Week (II) among Persons Aged 12 to 17: Percentages, 2002

| (I) Demographic Characteristic | Use LSD Once or Twice a Week ¹ | (II) Drug Use | Use LSD Once or Twice a Week ¹ | |
|---|---|--|---|-------------------------|
| | Great Risk | | Great Risk | Moderate/Slight/No Risk |
| TOTAL | 76.2 | PAST MONTH | | |
| AGE | | Any Illicit Drug ² | 11.2 | 13.6 |
| 12-13 | 69.7 | Marijuana | 8.1 | 9.0 |
| 14-15 | 76.7 | Any Illicit Drug Other Than Marijuana ² | 5.2 | 7.9 |
| 16-17 | 82.2 | Cigarettes | 13.2 | 12.6 |
| GENDER | | Alcohol | 18.3 | 16.5 |
| Male | 75.2 | Binge Alcohol ³ | 10.8 | 11.0 |
| Female | 77.3 | PAST YEAR | | |
| HISPANIC ORIGIN AND RACE | | Any Illicit Drug ² | 22.0 | 23.6 |
| Not Hispanic or Latino | 77.1 | Marijuana | 16.1 | 15.5 |
| White | 79.4 | Any Illicit Drug Other Than Marijuana ² | 12.7 | 16.8 |
| Black or African American | 69.9 | Cigarettes | 20.8 | 19.7 |
| American Indian or Alaska Native | 71.1 | Alcohol | 36.3 | 30.8 |
| Native Hawaiian or Other Pacific Islander | * | LIFETIME | | |
| Asian | 69.2 | Any Illicit Drug ² | 30.6 | 32.1 |
| Two or More Races | 75.1 | Marijuana | 21.4 | 19.2 |
| Hispanic or Latino | 71.5 | Any Illicit Drug Other Than Marijuana ² | 20.3 | 25.1 |
| COUNTY TYPE | | Cigarettes | 34.1 | 31.7 |
| Large Metro | 75.5 | Alcohol | 45.3 | 39.1 |
| Small Metro | 76.4 | | | |
| Nonmetro | 77.9 | | | |

*Low precision; no estimate reported.

¹ Respondents with missing data were excluded.

² Any Illicit Drug includes marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically. Any Illicit Drug Other Than Marijuana includes cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002.

Table 3.11A Perceived Great Risk of Having 4 or 5 Drinks of an Alcoholic Beverage Nearly Every Day, by Demographic Characteristics (I), and Selected Drug Use, by Risk Perceptions of Having 4 or 5 Drinks of an Alcoholic Beverage Nearly Every Day (II) among Persons Aged 12 to 17: Numbers in Thousands, 2002

| (I) Demographic Characteristic | Have 4 or 5 Drinks of Alcoholic Beverage Nearly Every Day ¹ | (II) Drug Use | Have 4 or 5 Drinks of Alcoholic Beverage Nearly Every Day ¹ | |
|---|--|--|--|-------------------------|
| | Great Risk | | Great Risk | Moderate/Slight/No Risk |
| TOTAL | 15,288 | PAST MONTH | | |
| AGE | | Any Illicit Drug ² | 1,239 | 1,627 |
| 12-13 | 5,317 | Marijuana | 859 | 1,157 |
| 14-15 | 5,183 | Any Illicit Drug Other Than Marijuana ² | 593 | 822 |
| 16-17 | 4,788 | Cigarettes | 1,379 | 1,813 |
| GENDER | | Alcohol | 1,945 | 2,403 |
| Male | 7,142 | Binge Alcohol ³ | 989 | 1,645 |
| Female | 8,146 | PAST YEAR | | |
| HISPANIC ORIGIN AND RACE | | Any Illicit Drug ² | 2,596 | 2,884 |
| Not Hispanic or Latino | 12,873 | Marijuana | 1,769 | 2,127 |
| White | 9,391 | Any Illicit Drug Other Than Marijuana ² | 1,528 | 1,802 |
| Black or African American | 2,339 | Cigarettes | 2,287 | 2,732 |
| American Indian or Alaska Native | 104 | Alcohol | 4,300 | 4,229 |
| Native Hawaiian or Other Pacific Islander | * | LIFETIME | | |
| Asian | 741 | Any Illicit Drug ² | 3,769 | 3,844 |
| Two or More Races | 255 | Marijuana | 2,387 | 2,702 |
| Hispanic or Latino | 2,415 | Any Illicit Drug Other Than Marijuana ² | 2,565 | 2,703 |
| COUNTY TYPE | | Cigarettes | 4,056 | 4,149 |
| Large Metro | 7,992 | Alcohol | 5,615 | 5,093 |
| Small Metro | 4,462 | | | |
| Nonmetro | 2,834 | | | |

*Low precision; no estimate reported.

¹ Respondents with missing data were excluded.

² Any Illicit Drug includes marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically. Any Illicit Drug Other Than Marijuana includes cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002.

Table 3.11B Perceived Great Risk of Having 4 or 5 Drinks of an Alcoholic Beverage Nearly Every Day, by Demographic Characteristics (I), and Selected Drug Use, by Risk Perceptions of Having 4 or 5 Drinks of an Alcoholic Beverage Nearly Every Day (II) among Persons Aged 12 to 17: Percentages, 2002

| (I) Demographic Characteristic | Have 4 or 5 Drinks of Alcoholic Beverage Nearly Every Day ¹ | (II) Drug Use | Have 4 or 5 Drinks of Alcoholic Beverage Nearly Every Day ¹ | |
|---|--|--|--|-------------------------|
| | Great Risk | | Great Risk | Moderate/Slight/No Risk |
| TOTAL | 62.2 | PAST MONTH | | |
| AGE | | Any Illicit Drug ² | 8.1 | 17.5 |
| 12-13 | 64.9 | Marijuana | 5.6 | 12.4 |
| 14-15 | 61.6 | Any Illicit Drug Other Than Marijuana ² | 3.9 | 8.8 |
| 16-17 | 60.0 | Cigarettes | 9.0 | 19.5 |
| GENDER | | Alcohol | 12.7 | 25.8 |
| Male | 56.9 | Binge Alcohol ³ | 6.5 | 17.7 |
| Female | 67.7 | PAST YEAR | | |
| HISPANIC ORIGIN AND RACE | | Any Illicit Drug ² | 17.0 | 31.0 |
| Not Hispanic or Latino | 62.2 | Marijuana | 11.6 | 22.9 |
| White | 60.7 | Any Illicit Drug Other Than Marijuana ² | 10.0 | 19.4 |
| Black or African American | 65.3 | Cigarettes | 15.0 | 29.4 |
| American Indian or Alaska Native | 55.7 | Alcohol | 28.1 | 45.5 |
| Native Hawaiian or Other Pacific Islander | * | LIFETIME | | |
| Asian | 74.5 | Any Illicit Drug ² | 24.7 | 41.3 |
| Two or More Races | 65.3 | Marijuana | 15.6 | 29.1 |
| Hispanic or Latino | 62.0 | Any Illicit Drug Other Than Marijuana ² | 16.8 | 29.1 |
| COUNTY TYPE | | Cigarettes | 26.5 | 44.6 |
| Large Metro | 64.8 | Alcohol | 36.7 | 54.8 |
| Small Metro | 60.9 | | | |
| Nonmetro | 57.5 | | | |

*Low precision; no estimate reported.

¹ Respondents with missing data were excluded.

² Any Illicit Drug includes marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically. Any Illicit Drug Other Than Marijuana includes cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002.

Table 3.12A Perceived Great Risk of Having 5 or More Drinks of an Alcoholic Beverage Once or Twice a Week, by Demographic Characteristics (I), and Selected Drug Use, by Risk Perceptions of Having 5 or More Drinks of an Alcoholic Beverage Once or Twice a Week (II) among Persons Aged 12 to 17: Numbers in Thousands, 2002

| (I) Demographic Characteristic | Have 5 or More Drinks of Alcoholic Beverage Once or Twice a Week ¹ | (II) Drug Use | Have 5 or More Drinks of Alcoholic Beverage Once or Twice a Week ¹ | |
|---|---|--|---|-------------------------|
| | Great Risk | | Great Risk | Moderate/Slight/No Risk |
| TOTAL | 9,379 | PAST MONTH | | |
| AGE | | Any Illicit Drug ² | 783 | 2,079 |
| 12-13 | 3,365 | Marijuana | 501 | 1,512 |
| 14-15 | 3,136 | Any Illicit Drug Other Than Marijuana ² | 394 | 1,021 |
| 16-17 | 2,879 | Cigarettes | 887 | 2,302 |
| GENDER | | Alcohol | 1,088 | 3,259 |
| Male | 4,287 | Binge Alcohol ³ | 535 | 2,100 |
| Female | 5,092 | PAST YEAR | | |
| HISPANIC ORIGIN AND RACE | | Any Illicit Drug ² | 1,515 | 3,954 |
| Not Hispanic or Latino | 7,797 | Marijuana | 999 | 2,892 |
| White | 5,349 | Any Illicit Drug Other Than Marijuana ² | 874 | 2,449 |
| Black or African American | 1,769 | Cigarettes | 1,435 | 3,575 |
| American Indian or Alaska Native | 61 | Alcohol | 2,468 | 6,051 |
| Native Hawaiian or Other Pacific Islander | * | LIFETIME | | |
| Asian | 440 | Any Illicit Drug ² | 2,306 | 5,287 |
| Two or More Races | 149 | Marijuana | 1,417 | 3,665 |
| Hispanic or Latino | 1,582 | Any Illicit Drug Other Than Marijuana ² | 1,567 | 3,685 |
| COUNTY TYPE | | Cigarettes | 2,526 | 5,665 |
| Large Metro | 4,790 | Alcohol | 3,373 | 7,320 |
| Small Metro | 2,768 | | | |
| Nonmetro | 1,821 | | | |

*Low precision; no estimate reported.

¹ Respondents with missing data were excluded.

² Any Illicit Drug includes marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically. Any Illicit Drug Other Than Marijuana includes cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002.

Table 3.12B Perceived Great Risk of Having 5 or More Drinks of an Alcoholic Beverage Once or Twice a Week, by Demographic Characteristics (I), and Selected Drug Use, by Risk Perceptions of Having 5 or More Drinks of an Alcoholic Beverage Once or Twice a Week (II) among Persons Aged 12 to 17: Percentages, 2002

| (I) Demographic Characteristic | Have 5 or More Drinks of Alcoholic Beverage Once or Twice a Week ¹ | (II) Drug Use | Have 5 or More Drinks of Alcoholic Beverage Once or Twice a Week ¹ | |
|---|---|--|---|-------------------------|
| | Great Risk | | Great Risk | Moderate/Slight/No Risk |
| TOTAL | 38.2 | PAST MONTH | | |
| AGE | | Any Illicit Drug ² | 8.3 | 13.7 |
| 12-13 | 41.2 | Marijuana | 5.3 | 10.0 |
| 14-15 | 37.4 | Any Illicit Drug Other Than Marijuana ² | 4.2 | 6.7 |
| 16-17 | 36.1 | Cigarettes | 9.5 | 15.2 |
| GENDER | | Alcohol | 11.6 | 21.5 |
| Male | 34.2 | Binge Alcohol ³ | 5.7 | 13.8 |
| Female | 42.4 | PAST YEAR | | |
| HISPANIC ORIGIN AND RACE | | Any Illicit Drug ² | 16.2 | 26.1 |
| Not Hispanic or Latino | 37.7 | Marijuana | 10.7 | 19.1 |
| White | 34.7 | Any Illicit Drug Other Than Marijuana ² | 9.3 | 16.1 |
| Black or African American | 49.5 | Cigarettes | 15.3 | 23.6 |
| American Indian or Alaska Native | 32.4 | Alcohol | 26.3 | 39.9 |
| Native Hawaiian or Other Pacific Islander | * | LIFETIME | | |
| Asian | 44.3 | Any Illicit Drug ² | 24.6 | 34.9 |
| Two or More Races | 38.2 | Marijuana | 15.1 | 24.2 |
| Hispanic or Latino | 40.7 | Any Illicit Drug Other Than Marijuana ² | 16.7 | 24.3 |
| COUNTY TYPE | | Cigarettes | 26.9 | 37.4 |
| Large Metro | 38.9 | Alcohol | 36.0 | 48.3 |
| Small Metro | 37.8 | | | |
| Nonmetro | 37.0 | | | |

*Low precision; no estimate reported.

¹ Respondents with missing data were excluded.

² Any Illicit Drug includes marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically. Any Illicit Drug Other Than Marijuana includes cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002.